February 22, 2022

Nemours Children's Health 10140 Centurion Parkway North Jacksonville, FL 32256

The Honorable Ron Wyden Chairman appropriate supports. Unfortunately, it is estimated that m ore than 45% of children diagnosed with a behavioral health disorder do not receive treatment.

At Nemours Children's H ospital, Delaware, our emergency department saw an increase of more than 80% in visits for suicidality or intentional harmin 2021 compared to 2020. Nemours Children's H ospital, Florida from 2020 to 2021, saw a 55% increase in patients in our emergency department with chief concerns of suicidality or intentional harm. Our behavioral health providers across our system have shared that our patients are increasingly experiencing higher levels of anxiety and depression, and grief from deaths of caregivers or family members. In outpatient and ambulatory care across our Florida operations, 85% of children screened had anxiety, depression, or another form of a behavioral health symptom.

We applaud the Surgeon General for raising the youth mental health crisis as a priority public health challenge. As the Surgeon General notes in his advisory, it will take time to resolve the many mental, emotional and behavioral (MEB) health challenges that children and youth are facing. However, the time to begin is now. We urge Congress to consider these five priorities to address barriers to providing high quality pediatric and youth mental health preventive services, supports and care:

- Address the social factors that contribute to poor mental health
- Support the pediatric MEB health workforce
- Strengthen reimbursement for MEB health services
- Sustain and expand access (testa) Tal 300 8 et 41.8 (s) 422 (5 of t10 T3 (e) .12) 374/T0 T2 L15 Ts (0) .462 v0 3 T of 20 3.5 of i t of 5,174 (c) (1,00 5 T

and test integrated, community based pediatric collaborations that align financial incentives and resources across Medicaid and other public and private programs to address SDOH improve MEB health and well-being, and reduce health disparities among pediatric populations. Models would be designed with input and engagement from community residents, Medicaid beneficiaries, and organizations, and be informed by a comprehensive needs and assets assessment in target communities.

Additionally, we encourage Congress to d irect CMS to review the early and periodic, screening, diagnostic and treatment (EPSDT) requirements and how they are being implemented across the states to support access to needed mental health services and early immost limber limbe

ⁱ Centers for Disease Control and Prevention. (2020). COVID -19 parental resources kit . Retrieved 2021, May 11 from https://www.cdc.gov/coronavirus/2019 -ncov/daily -life -coping/parental -resource - kit/index.html

vi

ⁱⁱ Brown, S. M., Orsi, R., Chen, P. C. B., Everson, C. L., & Fluke, J. (2021). The impact of the COVID -19 pandemic on child protection system referrals and responses in Colorado, USA. Child maltreatment 10775595211012476.

Leeb, R. T., Bitsko, R. H., Radhakrishnan, L., Martinez, P., Njai, R., & Holland, K. M. (2020). Mental health-related emergency department visits among children aged <18 years during the COVID -19

Pandemic - United States, January 1 -October 17, 2020. MMWR. Morbidity and mortality weekly report , 69(45), 1675–1680. https://doi.org/10.15585/mmwr.mm6945a3

Krass, P., Dalton, E., Doupnik, S. K., & Esposito, J. (2021). US pediatric emergency department visits for mental health conditions during the COVID -19 Pandemic. JAMA Network Open, 4 (4), e218533-e218533. https://doi.org/10.1001/jamanetworkopen.2021.8533

^v Centers for Disease Control and Prevention. (2021, March 22). Data and statistics on children's mental health. Retrieved 2021, May 27 from https://www.cdc.gov/childrensmentalhealth/data.html